Huntington Disease 101 – From a Clinician's and Family Member's Perspective

Presenters:


- Carol Eckert, Retired from Toronto Board of Education and mother of a son with Huntington's disease in a long term care facility
- Corey Janke, M.Ed, BSW, RSW, Southwestern Ontario Resource Center Director, Huntington Society of Canada



www.huntingtonsociety.ca

Carol's Story









Huntington Disease

- An inherited, degenerative, neuropsychiatric disease
- Characterized by a progressive loss of control over movements, thinking and emotions



HD - Some Numbers

- There is a 50% chance of inheriting the HD gene from the gene-carrying parent
- Approx. 1:7000 People in Canada have HD
- Average age of onset 35-50 years of age
- Average life span after dx: 15-20 years





HD - Genetics

- 46 Chromosomes 23 each parent
- chromosomes made up of genes
- genes made up of DNA
- DNA is made up of bases: Adenine, Thymine, Guanine, Cytosine
- HD gene located on chromosome 4
- Characterized by an expanded CAG repeat



Physical Impairments

Clumsiness/falling

Unsteady gait

Involuntary dance-like movements

Slurred speech

Difficulty swallowing

Intoxicated appearance

Difficulty sleeping

Excessive perspiring/body temperature

Locked-in" (late stages)



Cognitive Impairments

Memory

Impulse control

Slowed information processing

Decision making

Problem solving

Confusion

Judgement

Rigid (stubborn)



Selective Cognitive Changes

Impaired

Memory Retrieval

Verbal Fluency

Word Finding

Intact


Recognition Memory

Long-term Memory

Language Comprehension



Emotional Disturbances





Emotional/Behavioural Impairments

Short Temper

Depression

Apathy

Irritability

Verbal Outbursts

Assaultive

Psychosis (rare)

Mood Swings

Suicide



Resulting Emotional Behaviour

Apathy

Impulsiveness

Irritability

Social Withdrawal

Obsessive Compulsive Behaviours

Depression

Anxiety

Sleep Reversal or insomnia Suicidal Thoughts or Acts



Psychiatric and Behavioural Changes in HD

- Over 33% of HD persons develop psychiatric symptoms as the initial manifestation of illness
- Up to 73% of HD persons have concomitant psychiatric diagnoses-
- > 50% with affective disorder
- ≥ 10% with psychotic disorder
- Unknown % -intermittent explosive disorder



For More Information Contact:

Corey Janke
cjanke@huntingtonsociety.ca
Director of SWO Resource Centre
Huntington Society of Canada
(519) 660-0670